



ISSN: 2707-1146
e-ISSN: 2709-4189

Nature & Science

International Scientific Journal



Nature & Science

International Scientific Journal

Volume: 7 Issue: 12

2025

International Indices

ISSN: 2707-1146
e-ISSN: 2709-4189
DOI: 10.36719



Scilit



© It is necessary to use reference while using the journal materials.
© nature.science2000@aem.az
© info@aem.az
© aem.az

Founder and Editor-in-Chief

Researcher Mubariz HUSEYINOV, Azerbaijan Science Center / Azerbaijan
<https://orcid.org/0000-0002-5274-0356>
tedqiqat1868@gmail.com

Editor

Assoc. Prof. Dr. Akif AGHBABALI, Baku State University / Azerbaijan
<https://orcid.org/0000-0001-5793-9252>
oruclu@rambler.ru

Assistant editors

Assoc. Prof. Dr. Fuad RZAYEV, Institute of Zoology of MSERA / Azerbaijan
<https://orcid.org/0000-0002-8128-1101>
fuad.rzayev01f@gmail.com

Assoc. Prof. Dr. Saliga GAZI, Institute of Zoology of MSERA / Azerbaijan
<https://orcid.org/0000-0002-9378-4283>
seliqegazi08@gmail.com

Researcher Gulnar ALIYEVA, Azerbaijan Science Center / Azerbaijan
<https://orcid.org/0009-0004-1769-777X>
gulnar.musayeva1982@gmail.com

Language editor

Assoc. Prof. Dr. Gunay RAFIBAYLI, Baku State University / Azerbaijan

Editors in scientific fields

Prof. Dr. Ibrahim MAMMADOV, Baku State University / Azerbaijan
Assoc. Prof. Abulfaz TAGHIYEV, Baku State University / Azerbaijan

EDITORIAL BOARD

Prof. Dr. Aziz SANJAR, University of North Carolina at Chapel Hill / USA
Prof. Dr. Irada HUSEYNOVA, Institute of Molecular Biology and Biotechnology of MSERA / Azerbaijan
Prof. Dr. Vagif ABBASOV, Institute of Petrochemical Processes named after Academician Y. H. Mammadaliyev of MSERA / Azerbaijan
Prof. Dr. Nazim MURADOV, University of Central Florida / USA
Prof. Dr. Vagif FARZALIYEV, Institute of Chemistry of Additives named after M. Guliyev of the MSERA / Azerbaijan
Prof. Dr. Georgi DUKA, Moldovan Academy of Sciences / Moldova
Prof. Dr. Ibrahim JAFAROV, Scientific Research Institute of Plant Protection and Technical Plants of the MARA / Azerbaijan
Prof. Dr. Elshad GURBANOV, Baku State University / Azerbaijan
Prof. Dr. Aliaddin ABBASOV, Nakhchivan State University / Azerbaijan
Prof. Dr. Ilham SHAHMURADOV, Institute of Genetic Resources of MSERA / Azerbaijan
Prof. Dr. Ulduz HASHIMOVA, Institute of Physiology of MSERA / Azerbaijan
Prof. Dr. Aloysat GULIYEV, Institute of Soil Science and Agro Chemistry of MSERA / Azerbaijan
Prof. Dr. Sayyara IBADULLAYEVA, Institute of Botany of MSERA / Azerbaijan
Prof. Dr. Ismayıl ALIYEV, Baku State University / Azerbaijan
Prof. Dr. Dunya BABANLI, Azerbaijan State Oil and Industry University / Azerbaijan
Prof. Dr. Mehmet KARATASH, Nejmettin Erbakan University / Turkey
Prof. Dr. Shaig IBRAHIMOV, Institute of Zoology of MSERA / Azerbaijan
Prof. Dr. Shahlar BABAYEV, Azerbaijan State Agrarian University / Azerbaijan
Prof. Dr. Afat MAMMADOVA, Baku State University / Azerbaijan
Prof. Dr. Akbar AGHAYEV, Sumgayit Statet University / Azerbaijan
Prof. Dr. Rajes KUMAR, Ministry of Textile / India
Prof. Dr. Bahadur GALANDAROV, Baku State University / Azerbaijan
Prof. Ali AZGANI, University of Texas at Taylor / USA
Assoc. Prof. Dr. Rinat ISKAKOV, Atyrau Oil and Gas University / Kazakhstan
Assoc. Prof. Dr. Elshad ABDULLAYEV, Sumgayit Statet University / Azerbaijan
Assoc. Prof. Dr. Bushra BILAL, Muhammad Ali Jinnah University / Pakistan
Assoc. Prof. Dr. Alia RZAYEVA, Nakhchivan State University / Azerbaijan
Assoc. Prof. Dr. Elchin HUSEYN, Rotterdam Erasmus University Medical Center / Netherlands
Dr. Svetlana GORNOVSKAYA, Beloserkovsk National Agrarian University / Ukraine

Metabolism of the Bilirubin and its Biochemistry Role

Nurgun Mehbaliyeva¹ , Narmina Abdullayeva^{2*} , Nigar Huseynova² 

Abstract. Bilirubin is a tetrapyrrolic bile pigment that plays a crucial role in several vital metabolic pathways of the human body. It is a yellow-colored compound that circulates in the bloodstream and participates in various physiological processes. Bilirubin is predominantly generated through the catabolism of hemoglobin following the breakdown of erythrocytes, and its biosynthesis is primarily mediated by the liver. The homeostasis of bilirubin is intricately linked to multiple physiological and biochemical functions, and its precise regulation is essential for maintaining systemic health. Elevated serum bilirubin levels may serve as a clinical biomarker for a range of pathological conditions, including hepatic dysfunctions, hematological disorders, and cholestatic syndromes.

Keywords: bilirubin, free radicals, liver, Kupffer cells, cholesterol

Introduction

Bilirubin metabolism affects various physiological and biochemical processes of the body, and its proper regulation is essential for maintaining health. Bilirubin exists in two major forms: conjugated and unconjugated bilirubin. Direct bilirubin is generated by the conjugation process in the liver and excreted from the body by being excreted as waste in the moist intestines. Conjugated bilirubin is formed in the liver through the conjugation process and is excreted into the intestines as a waste product, eventually being eliminated from the body. Unconjugated bilirubin, on the other hand, is produced as a result of erythrocyte degradation and is processed in the liver where it is converted into direct (conjugated) bilirubin. Disruptions in this metabolic pathway can lead to hyperbilirubinemia, a condition clinically manifested as jaundice. Notably, bilirubin is not merely a metabolic waste product; it also functions as a physiological antioxidant, helping to neutralize harmful free radicals in the body (Chen et al., 2018).

Materials and Methods

Thus, the liver serves as the central organ in the synthesis, excretion, and metabolism of bilirubin. Bilirubin originates from the degradation of erythrocytes, which have a lifespan of approximately 120 days, and its proper handling is essential for maintaining liver health. Hepatic metabolic processes—particularly the production and excretion of bilirubin—constitute key aspects of this function. The presence and concentration of bilirubin serve as important biomarkers for the early detection of liver and hematological disorders, providing a valuable basis for future clinical and molecular research (Agius, 2018).

¹ Baku State University, Bachelor student, Baku, Azerbaijan

² Baku State University, PhD in Biology, Baku, Azerbaijan

*Corresponding author. E-mail: narminaabdullayeva@bsu.edu.az

Received: 19 September 2025; Accepted: 20 November 2025; Published online: 25 December 2025

© The Author(s) 2025. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Hepatic Glucose Metabolism. Hepatocytes are the primary cell type involved in glucose metabolism within the liver. Circulating glucose enters hepatocytes via the GLUT2 transporter, which is embedded in the plasma membrane. The specific deletion of GLUT2 in hepatocytes abolishes hepatic glucose uptake. In addition, GLUT2 facilitates the release of glucose from the liver into the bloodstream. However, the absence of GLUT2 does not impair hepatic glucose production during fasting, suggesting that glucose may also be released via alternative transporters (e.g., GLUT1) or other mechanisms (Evans et al., 2013). Once inside the hepatocyte, glucose is phosphorylated by the enzyme glucokinase into glucose-6-phosphate (G6P). This reaction reduces intracellular free glucose concentrations, thereby facilitating continued glucose uptake. Since G6P cannot be transported out of the cell, it remains within hepatocytes. In the fed state, G6P serves as a key precursor for glycogen synthesis. Additionally, G6P is metabolized to pyruvate via glycolysis. Pyruvate then enters the mitochondria and is fully oxidized through the tricarboxylic acid (TCA) cycle and oxidative phosphorylation, leading to ATP production (Fig. 1).




Figure 1. Glycogen metabolism

In the fed state, glucose enters hepatocytes via GLUT2, is phosphorylated by glucokinase, and subsequently utilized by glycogen synthase for glycogen synthesis. During fasting, glycogen is hydrolyzed by glycogen phosphorylase to release glucose through glycogenolysis (Chew et al., 2023). The liver is considered the central regulatory organ responsible for maintaining whole-body cholesterol homeostasis (Fig. 2).




Figure 2. Structure of cholesterol

It serves as the primary site for de novo cholesterol biosynthesis, facilitates the clearance of cholesterol-containing chylomicron remnants and low-density lipoprotein (LDL) particles from the plasma, and makes a major contribution to the formation of high-density lipoproteins (HDL) (Zhang et al., 2023). Approximately 4 mg/kg of bilirubin is produced daily. Heme is a tetrapyrrolic macrocycle consisting of four pyrrole rings interconnected by carbon bridges, with a central iron atom. Bilirubin is primarily generated through a two-step sequential enzymatic degradation of heme within the reticuloendothelial system, particularly in the spleen. Other contributing cells include phagocytes and Kupffer cells of the liver. Once released into the plasma, bilirubin binds to albumin, the main transport protein in the bloodstream (Pirone et al., 2009).

The active transport of unconjugated bilirubin is mediated by carrier proteins, although the exact identity and mechanism of these transporters remain poorly understood (Fig. 3) (Blumgart et al., 2017). Albumin has a very high binding affinity for bilirubin, and under physiological conditions, unbound (free) unconjugated bilirubin is virtually undetectable in the plasma (Chew et al., 2023). Bilirubin is taken up by hepatocytes from the hepatic sinusoids via two distinct mechanisms: passive diffusion and receptor-mediated endocytosis. Passive diffusion does not require energy and occurs along the concentration gradient, resulting in a bidirectional flow.




Figure 3. Bilirubin synthesis in the liver

Results and Discussion

The study demonstrated, that bilirubin is an important metabolite of heme (ferroprotoporphyrin IX), a coordination complex that links iron in various proteins. It is a potentially toxic substance. However, the body has developed mechanisms for its safe detoxification and excretion. Bilirubin and its metabolites also provide a distinctive yellow color of bile and feces and, to a lesser extent, urine. This topic summarizes the mechanism of heme metabolism and bilirubin synthesis (Bauer & Kämper, 2016). Bilirubin is formed by a 2-stage sequential catalytic degradation reaction that occurs primarily in the cells of the reticuloendothelial system, specifically in the spleen. Other cells include phagocytes and Kupfer cells of the liver. Receiving Heme these cells receive heme and the enzyme heme oxygenase acts on them. The enzyme releases chelated iron by catalyzing the oxidation of the alpha-carbon bridge. This reaction produces equimolar amounts of carbon monoxide, which is excreted by the lungs and leads to the formation of the green pigment biliverdin (Chisari & Ferrari, 2018). Bilirubin, insoluble in aqueous solution, binds in circulation to albumin, which is a reversible and covalent type of binding.

Metabolism of bilirubin. Albumin binding: after bilirubin is released into plasma, it is taken up by albumin, a carrier throughout the body. The binding affinity of albumin to bilirubin is quite high, and under ideal conditions, free (non-albumin-bound) unconjugated bilirubin does not appear in plasma. To a lesser extent, especially in cases of hypoalbuminemia, binding with high-density lipoproteins

also occurs. Albumin binding limits the outflow of bilirubin from the vascular cavity, minimizes glomerular filtration and prevents its deposition and deposition in tissues (de Sauvage et al., 2011). Bilirubin is taken from hepatic sinusoids into hepatocytes by two different mechanisms: passive diffusion and Receptor-Mediated Endocytosis. The passive diffusion process does not consume energy and, as a result, proceeds with a concentration gradient, making the flow bidirectional. Active transporter intake of unconjugated bilirubin from hepatic sinusoids is carried out through carrier proteins that are not well understood. Part of the conjugated and unconjugated bilirubin within the hepatocyte is transported back into the sinusoidal space, and this fraction is again transported downstream of the sinusoidal flow. Conjugation is mandatory to dissolve bilirubin in water and facilitate its secretion through the canalicular membrane and excretion with bile. Bilirubin binds to glucuronic acid in the hepatocyte by a family of enzymes called uridine-diphosphoglucuronic glucuronosyltransferase (UDPGT) (Sedlak et al., 2017). The process of glucuronidation is one of the many important detoxification mechanisms of the human body. Under normal conditions, bilirubin is the main molecule from which diglucuronide is synthesized. However, if the conjugation system fails under conditions of excessive bilirubin synthesis, most of the bilirubin may be conjugated as bilirubin monoglucuronide. The combination of bilirubin into a water-soluble form involves breaking hydrogen bonds, an important process for its excretion by the liver and kidneys. This is achieved by glucuronic acid, which binds the side chains of propionic acid of bilirubin (Liu et al., 2008).

Conclusion

In summary, the liver is the central organ involved in the synthesis, excretion, and metabolism of bilirubin. Bilirubin is generated from the degradation of erythrocytes, which have a lifespan of approximately 120 days, and its proper processing is essential for maintaining hepatic function. The metabolic activities of the liver - particularly those related to bilirubin production and elimination - constitute critical aspects of its physiological role. The plasma concentration of bilirubin serves as a valuable biomarker for the early detection of both hepatic and hematologic disorders. Bilirubin metabolism affects various biochemical processes, and its regulation is essential for maintaining health, and it provides a foundation for diagnostic research.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Agius, L. (2018). Molecular aspects of glucokinase and hepatic glycogen metabolism. *Journal of Clinical Biochemistry & Metabolism*.
2. Bauer, C. P., & Kämper, C. (2016). The molecular mechanisms of bilirubin conjugation and transport. *European Journal of Clinical Investigation*.
3. Blumgart, L., Belghiti, J., Jarnagin, W., & DeMatteo, R. (Eds.) (2017). Chapter 7 - Biliary Tract Pathophysiology. *Surgery of the Liver, Biliary Tract and Pancreas* (4th ed.). W.B. Saunders.
4. Chisari, F., & Ferrari, C. (2018). The liver and its role in bilirubin metabolism. *Gastroenterology Clinics of North American*.
5. Chew, D., Bartola, S., & Schenck, P. (2023). Chapter 1 - Urinalysis. *Canine and Feline Nephrology and Urology* (2nd ed.). W.B. Saunders.
6. Chen, W., Maghzal, G., Ayer, A., Suarna, C., Dunn, L., & Stocker, R. (2018). Absence of the biliverdin reductase-a gene is associated with increased endogenous oxidative stress. *Journal of Radical Biology & Medicine*.
7. Evans, J., Morris, L., & Marchesi, J. (2013). The gut microbiome: The role of the virtual organ in host endocrinology. *Journal of Endocrinology*.

8. de Sauvage Nolting, P., Kusters, D., Hutten, B., & Kastelein, J. (2011). ExPRESS study group. Serum bilirubin levels in familial hypercholesterolemia: a new risk marker for cardiovascular disease. *Journal of Lipid Research*.
9. Sedlak, T., Saleh, M., Higginson, D., Paul, B., Juluri, K., & Snyder, S. (2017). Bilirubin and glutathione have complementary antioxidant and cytoprotective roles. *Proceedings of the National Academy of Sciences of the United States of America*.
10. Liu, Li., Lu, J., Xiong, W., Oger, J., Tetzlaff, W., & Cynader, M. (2008). Bilirubin possesses powerful immunomodulatory activity and suppresses experimental autoimmune encephalomyelitis. *Journal of Immunology*.
11. Pirone, C., Quirke, J., Martin, E., Horacio, A., & Lee, D. (2009). Animal Pigment Bilirubin Discovered in Plants. *Journal of the American Chemical Societ*.
12. Zhang, Y., Lai, Q., Chen, W., Zhang, C., & Liu, Z. (2023). Recent advance in cortisol immunosensing technologies and devices. *Chemosensors*.

Study of Diagnostic Methods and Control Measures For Listeriosis Disease Among Big-Horned Animals in the Ganja-Gazakh Zone

Ayten Agayeva^{1*} , Rahila Farmanli¹ 

Abstract. *The purpose of the research is to study the diagnostic methods and control measures for listeriosis among cattle in farms of the Ganja-Gazakh zone. The research was conducted in various farms located in Shamkir, Samukh and Goygol districts, in the laboratory of the Department of “Episootology, Microbiology and Parasitology” of the Faculty of “Veterinary Medicine” of the Azerbaijan State Agrarian University of Veterinary Medicine. Depending on the immune status of the organism and the virulence of the pathogen, the disease manifests itself in acute, and subacute forms among animals. Based on the high efficiency of their application in combination with sulfagin, gentomycin and oxytetracycline for the specific treatment of the disease, we recommend their use for the prevention of listeriosis before vaccination. In case of listeriosis, along with specific treatment, we carried out symptomatic treatment. We carried out disinfection, disinsection and deratization works in livestock farms in an organized manner.*

Keywords: *cattle, disease, listeriosis, bacteriological examination, morphological, cultural, biochemical, antigenic, pathogenic properties*

Introduction

Listeriosis is an infectious disease of animals and humans caused by the bacteria *Listeria monocytogenes* and is characterized by nervous system disorders, septic conditions, puerperal disease and mastitis. An asymptomatic (latent) form of the disease is also observed. Many species of domestic and wild mammals, birds and humans are affected. It can occur as a secondary or mixed infection in a number of infectious diseases in pigs and birds. The disease has been recorded in Azerbaijan (Shirinov, 2002; Eyubov, 2005).

Many scientists have determined that the occurrence of listeriosis in various types of animals and birds depends on the amount of precipitation, the nature of the feed, pH, the degree of contamination of the soil with the causative agent of the disease, the presence of antagonistic microflora in the soil and the degree of virulence of the causative agent of listeriosis. P. R. Lazarev, V. I. Gershun and I.I.Guslavskaya show that incompleteness of the feed ration due to cold and soluble protein, as well as changes in the pH of silage, reduce the overall resistance of animals and increase their susceptibility to listeriosis (Kalinin, 1987).

¹ Azerbaijan State Agrarian University, Baku, Azerbaijan

*Corresponding author. E-mail: aayeva83@mail.ru

Received: 6 September 2025; Accepted: 13 November 2025; Published online: 25 December 2025

© The Author(s) 2025. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Listeriosis causes a decrease in animal productivity, causes metritis during calving, and causes great economic damage. At the same time, it causes a certain amount of financial resources to be spent on various health measures (Mammadli, 2015).

Listeriosis is a zoonotic disease of animals and humans, characterized by polymorphism of clinical signs or asymptomatic carriage and high mortality. M.M. Halimbayov, A.A. Annaghiyev, M.G. Ganiyev, A.M. Khalilov, A. Khankishiyev, H.A. Rahimova, Y. Ahmadov, C.N. Mammadova, A.G. Abbasov, F.M. Gulubayov played a special role in studying the epizootology, pathogenesis and specific prophylaxis of listeriosis in various types of agricultural animals in the Republic of Azerbaijan. In our country, the first vaccine against listeriosis of agricultural animals in world veterinary practice was developed and this preparation was used in the specific prophylaxis of listeriosis of sheep in 1965–1978. In 1980, a colored antigen was prepared for the diagnosis of listeriosis of agricultural animals with AR (A.A. Annaghiyev, E.A. Aliyev, F.M. Gulubeyov), an author's certificate was obtained for the diagnostic, and it is currently used in both veterinary and medical practice (Aliyev, 2013; Mammadli, 2020).

Listeriosis is an infectious disease characterized by meningoencephalitis, balanitis, metritis, mastitis, septicemia. The incubation period lasts from one week to 1 month. Listeriosis occurs in septic, nervous, genital and atypical forms. Factor (0.3-0.5 and 0.5-2.0 μm) is a bacterium located singly, in pairs or in clusters. In some cases, bacteria occur in polymorphic rod, cocci, vibrio forms. Bacteria stain positively with all aniline dyes and the Gram method. Microbes do not form capsula and spores. In fresh (4-6 hours) broth culture, bacteria are motile. On meat-peptone agar-agar, bacteria form thin, fleshy colonies similar to *Pasteurella*, with a diameter of 1-1.5 mm (Lysak, 2007).

When studying culture strains obtained from various sources by I.A. Bakulov, O.V. Krivonosov, I.V. Gershun, it was determined that there are both virulent and weakly virulent variants of listeria. In addition, various literature data show that external environmental factors play an important role in the occurrence and course of listeriosis (Bessarabov, 2007).

The disease has a mortality rate of up to 40%, and causes considerable damage to agriculture as a result of reduced productivity and calving in animals. In addition, a system of various health measures is used, which leads to the expenditure of a certain amount of financial resources (Gadimov, 1990).

Materials and Methods

To achieve the set goal, we selected 26 sick animals related to the red desert and local breeds, divided them into 6 groups of 4 animals in each group, and conducted scientific research on them. The research work was carried out in various cattle farms of the Ganja-Gazakh zone (Shamkir, Samukh and Goygol), at the Department of Epizootology, Microbiology and Parasitology of the Azerbaijan State Agricultural University. In recent years, the climate in the Republic of Azerbaijan has changed dramatically, the amount of precipitation has decreased and drought has increased. As a result, the composition of feed in many regions has changed, its quality has deteriorated, and the number of insects and rodents has increased. As we know, the main source of infection is various wild and domestic animals, especially rodents, birds and insects, which carry the causative agent of the disease. The microbe enters the body through the mucous membranes (nose, conjunctiva, mouth) through water, feed, air, dust, etc. It is assumed that infection also occurs through blood-sucking insects, especially ticks.

As a result of all these factors, listeriosis was recorded among cattle in various farms of Shamkir, Samukh and Goygol regions. Based on the appeal of the heads of these farms, employees of the Department of Epizootology, Microbiology and Parasitology of the ASAU were engaged in studying the occurrence, course, treatment and many other issues of the disease.

It turned out that the farms mentioned were healthy due to listeriosis and had no contact with unhealthy farms. Listeriosis infection was recorded in August-October 2022. The disease was mainly observed in 18-28-month-old heifers, and infection was not detected in the causative bulls. The incubation period of the disease is 7-30 days. In addition to fever, general weakness, and loss of appetite, some animals showed signs of central nervous system dysfunction, including loss of balance, convulsions, severe nervous excitability, paralysis of some muscle groups, neck flexion, and conjunctivitis. In a group of sick animals, blindness, stomatitis, anemia of visible mucous membranes, and comatose state were noted.

Listeriosis mainly has an acute, subacute, and chronic course. Listeriosis occurs in several clinical forms: nervous, septic, mixed, latent-asymptomatic, with damage to the reproductive system (parturition, delayed ejaculation, endometritis, and metritis) and udder (mastitis). In large horned animals, the CNS is more affected. The disease begins with weakness and loss of appetite. Serous-mucous discharge occurs from the nasal cavity, and abundant mucus is secreted from the mouth. After 3-7 days, uncontrolled movements, convulsions, fits of excitement, paresis of individual muscle groups, loss of vision, conjunctivitis, stomatitis are observed. Body temperature rises at the beginning of the disease or remains within the normal range. Listeriosis in calves proceeds in the form of septicemia, sometimes accompanied by damage to the CNS.

Results and Discussion

From the research work we conducted, it became clear that the pathological-anatomical changes in large horned animals were mainly sharp changes in the brain, especially the engorgement of cerebral blood vessels, blood leakages were observed on the brain and cerebellum. In the cranial cavities and ventricles of the brain, a cloudy fluid mixed with pus was seen. To diagnose listeriosis, we took into account its epizootological characteristics, clinical signs and pathological-anatomical changes, used blood culture, bacteriological examination and serological reactions. While the animal was alive, we used the blood of the sick animal (5-10 ml), the discharged pupa, pupal discharge, pupal membrane, nasal passages, conjunctival fluid, and milk taken from the inflamed scrotum. We also sent the brain and spinal cord to the laboratory for examination.

We examined the pathological samples received in the laboratory based on the bacteriological scheme. Bacteriological examinations play an important role in the diagnosis of listeriosis. During microscopy, we prepared a smear from the pathological material and stained it using the Gram method. During microscopy, listeria were found singly or in pairs. In some cases, we determined that listeria were observed in coccus-like, diplobacteria, rod-shaped, and chain-like forms.

We obtained a pure culture of the factor by inoculating the blood of sick animals from the parenchymatous organs of dead animals into various nutrient media, prepared smears on glass slides, examined them under a microscope, and used biological tests. After obtaining a pure culture of the listeriosis pathogen, it was examined for its morphological, cultural, tinctorial, and biochemical properties, and its pathogenicity was determined. It turned out that the pathogen forms a precipitate in liquid nutrient media and, when shaken, rises up like a hair, resembling small, transparent dewdrops in solid nutrient media.

Table 1.
Treatment of cattle with listeriosis

Treatments	Injection method	Dose (once)	Number of patients	Conclusion	
				Survived quantity	%
Sulfagin and gentomisin	Perosal and intramuscular	25 g+2.5 ml	4	4	100
Sulfagin and oxytetracycline	Peros	25 g+2.5 ml	4	4	100
Gentomycin	Intramuscular	2.5 ml	4	3	75
Oxytetracycline	Intramuscular	2.5 ml	4	3	75
Sulfagin	Peros	25 g	4	2	50
Control	–	–	4	4	–

As can be seen from Table 1, while sulfagin in combination with gentomycin and oxytetracycline achieves 100% recovery, only 50-75% of animals recover from the use of these drugs alone. Based on the high efficiency of sulfagin in combination with gentomycin and oxytetracycline, we recommend their use for the prevention of listeriosis before vaccination. In the case of listeriosis, along with specific treatment, we carried out symptomatic treatment. The main goal of symptomatic treatment is to stimulate the functional activity of the cardiovascular system and the digestive system. In order to ensure the healthy and vigorous development of large-horned animals, and to prevent them from contracting listeriosis, we set ourselves the goal of timely implementation of preventive measures. One of such measures and the main one is timely and high-quality vaccination of animals with the appropriate vaccine.

According to researchers, individuals who have contracted listeriosis and recovered from it in natural conditions develop relative immunity. Precipitins, agglutinins, and complement-binding antibodies are detected in the blood serum of animals that have recovered from the disease naturally. However, the serum of convalescents does not have therapeutic properties. The first vaccine against the disease was prepared by A.A. Annagiyev from the “A” strain. A dry live vaccine prepared from the “AUF” strain was proposed in 1974 and is used based on the current instructions (Alasgarov, 2016).

We vaccinated all clinically healthy animals 10 days after treatment with a dry vaccine prepared from the AUF strain against listeriosis. We vaccinated 687 cattle on farms according to the instructions. We injected the vaccine intramuscularly twice with an interval of 10 days. We periodically sent samples to the laboratory to determine the quality indicators of feed and water. We applied the feed ration, which is important for feeding large horned animals. We implemented the solution of the following issues in disease prevention. In order to ensure the protection of healthy livestock farms from the disease due to listeriosis:

- We ensured that animals brought to farms were kept in quarantine for 30 days.
- We ensured the systematic elimination of rodents.
- We eliminated blood-sucking insects and ticks.
- Ensuring constant quality control of feeds (especially silage and mixed feed)
- We ensured the conduct of bacteriological examinations in accordance with the relevant requirements.
- In cases of abortion or stillbirth, we implemented a strict approach and sent pathological samples for examination.

We boiled milk from large horned animals (with a positive result in serological reactions) for 15 minutes. We banned the transportation of feed that had been in contact with sick animals. We carried out disinfection, disinfestation and deratization work in livestock farms in an organized manner. We

biothermally neutralized manure on the farm. We disinfected stables and farmyards. For this purpose, we used 3% sodium hydroxide, 2% chlorinated lime solution, and 6% creolin emulsion. We periodically carried out deratization measures on the farm. We lifted the restriction 2 months after the last patient recovered from the farm and after receiving a negative result during serological tests.

Conclusion

The results of the study showed that 100% recovery was achieved when sulfagin was used in combination with gentamicin or oxytetracycline for the treatment of sick animals in unhealthy farms due to listeriosis. During the research work we conducted, we vaccinated all animals susceptible to listeriosis in unhealthy farms with a dry vaccine prepared from the AUF strain according to the relevant instructions. For the prevention of listeriosis, animals should always be brought to the farm from healthy farms. We always monitored the quality of feed and fought against rodents. We used preventive measures when the disease was observed on the farm.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Alasgarov, Z. A. (2016). *Anthropozoonous diseases*. Science.
2. Aliyev, E. A., Azimov, I. M., Valiyev, U. M., & Safi, N. V. (2013). *Epizootology and infectious diseases*. UniPrint.
3. Bessarabov, B. F., Vashutin, E. S., & Voronin, E. C. (2007). *Infectious diseases of animals*. Colossus.
4. Eyubov, I. Z., Hajiyeve, Y. H., Shirinov, F. B., Ahmadov, C. E., & Mammadov, A. T. (2015). *Veterinary medicine (reference book)*. Azernashr.
5. Gadimov, R. A., Mammadov, I. B., & Julfayev, J. A. (1990). *Special epizootology*. Baku University Publishing House.
6. Kalinin, N. M. (1978). *Listeriosis of large-horned cattle*. Colossus.
7. Lysak, V. V. (2007). *Microbiology*. BGU.
8. Mammadli, A., Abdullayev, M., Hasanov, A., & Salmanov, R. (2020). *Veterinary and sanitary service against infectious diseases in livestock*. Teacher Publishing House.
9. Mammadli, A., Murtuzov, G., & Shahmarov, A. (2015). *Protection of agricultural animals from diseases*. Teacher Publishing House.
10. Shirinov, F. B. (2002). *Explanatory dictionary of epizootological terms*. Ganges.
11. Wesley, I. V. (2007). Listeriosis in animals. In E. T. Ryser & E. H. Marth (Eds.). *Listeria, listeriosis, and food safety* (3rd ed.). CRC Press.
12. Wiedmann, M., Arvik, T., Bruce, J. L., Neubauer, J., Del piero F., Smith, M. C., Hurley, J., Mohammed, H. O., & Batt, C. A. (1997). Investigation of a listeriosis epizootic in sheep in New York State. *Am. J. Vet. Res.*, 58, 733–737.

The Possibilities of Using the Nature of Semiconductors in Practical Life

Gunay Dadashzade 

Abstract. *The transformation of the global automotive market with the increase in the production of electric vehicles in recent years has created an incredible demand for the development of electronic components based on SiC and GaN semiconductors with a wide band gap in the world, without which it is impossible to imagine the electronics of modern and future hybrid and electric vehicles, communication systems, mobile devices and space technology. This demand has led to a sharp increase in investment in the development and development of new technologies for SiC and GaN wafers of larger diameters in order to reduce the cost of manufacturing electronic products and continue to displace classic silicon components. Every year, semiconductor devices based on SiC and GaN are penetrating deeper into our lives. The past 2021 allowed them to make an impressive breakthrough and lay the foundation for growth in their consumption in the next five years and beyond. The main driver of this growth is the automotive market, or more precisely, hybrid and electric cars.*

Keywords: *electric vehicle, components, battery, power, semiconductors*

Introduction

Semiconductors are materials that can conduct electricity, but to a limited extent. Their unique feature is that their electrical conductivity can be controlled and modified by introducing appropriate dopants or by changing external conditions, such as temperature, pressure, or an electric field. Under normal conditions, semiconductors act as insulators, but under certain circumstances, they can conduct electricity, making them indispensable in the production of electronic components.

Natural semiconductors, such as pure silicon, do not have sufficient electrical properties for use in modern electronic devices. To increase their ability to conduct electricity, doping is used, which involves introducing small amounts of other chemical elements into the semiconductor structure. Depending on the type of dopant, two main types of doped semiconductors can be distinguished:

An n-type semiconductor is formed by doping a semiconductor with elements with a large number of electrons, such as phosphorus or arsenic. The introduction of these impurities creates an excess of electrons that can move freely within the material, increasing its conductivity. A p-type semiconductor is formed by doping a semiconductor with elements with a smaller number of electrons, such as boron or aluminum. These impurities form so-called electron holes, which act as positive charge carriers and also help improve the material's conductivity.

Azerbaijan State Pedagogical College, Baku, Azerbaijan

E-mail: gunay_dadaszade@mail.ru

Received: 12 September 2025; Accepted: 28 November 2025; Published online: 25 December 2025

© The Author(s) 2025. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

The most common semiconductor materials are silicon (Si) and germanium (Ge), although many other compounds with semiconducting properties exist, such as gallium arsenide (GaAs) and indium phosphide (InP). These materials are used to manufacture semiconductor components such as diodes, transistors, and integrated circuits, which form the basis of modern technology. The share of hybrid and electric vehicle sales in the country reached 94.9%. Norway is actively building charging and service infrastructure, and numerous tax breaks for citizens buying electric vehicles are stimulating the transition away from internal combustion engine vehicles. Next-generation electric vehicles require power devices that can improve the efficiency of the vehicle (with a subsequent increase in the driving range) and the speed of battery charging. Figure 1 shows the key nodes of the vehicle where SiC and GaN electronic components can be used. SiC inverters have proven to be a key solution to meet these requirements. In addition to converting the input DC to AC, the inverter regulates the level of power supplied to the motor in accordance with driving needs (Challenges and Future Perspectives).

Methods

The progress of silicon carbide technology, in terms of increasing diameter, production volume, improving quality and decreasing cost of SiC, has reached the point where mass production of 150 mm wafers is based on the use of silicon carbide blanks, as shown in Fig. 1. The progress of silicon carbide technology, in terms of increasing diameter, production volume, improving quality and decreasing cost of SiC, has reached the point where mass production of 150mm wafers is based on the use of silicon carbide blanks, as shown in Fig. 1.



Figure 1. Production of 150 mm thick silicon carbide plate

Findings and Discussion

The role of the inverter is increasing as the electric vehicle industry gradually transitions from 400 to 800 V. The efficiency of transferring battery power to the motor in a traditional inverter is 97–98%, while the efficiency of a SiC inverter reaches 99%. Note that an increase in efficiency of one or two decimal places provides very significant benefits to the entire vehicle (Compound Semiconductor Quarterly Market Monitor, 2021).




Figure 2. Key nodes for the application of SiC and GaN-based electronic components in an electric vehicle

SiC inverters are ideal for these applications because they can withstand high voltages and temperatures, allowing all other components to be smaller. With 800 V batteries, the required current is reduced and smaller cables can be used, reducing costs, vehicle weight, and electrical system assembly time. This not only improves the range of EVs, but also their efficiency. Charging times for 800 V batteries can be reduced to one-fifth of the time required for 400 V batteries by using powerful SiC DC/DC converters. The former's high efficiency allows the amount of energy delivered to the batteries to be maximized during charging with negligible power losses (SiC and GaN: A Tale of Two Semiconductors, 2021).

The global growth of the BEV market, which meets modern standards for efficiency and CO₂ emissions, requires the use of new semiconductor technologies in the drive inverter. The supply voltage of the BEV inverter is in the range of 400–900 V, depending on the drive power, battery type and the presence of a step-up converter. Since the drive inverter controls the motor, its operating frequency is usually less than 20 kHz. The advantage of using higher frequencies here is only to move away from the audible range of audio noise. Therefore, the main losses of the inverter are conduction losses, especially at low BEV loads (SiC and GaN: A tale of two semiconductors, 2021).

Typically, the choice in such cases is a silicon IGBT, but its inherent saturation threshold voltage (due to its "bipolar" structure) at low loads cannot be reduced, even when a large number of IGBTs are connected in parallel. Silicon carbide has an electric field strength 10 times higher (~3 MV/cm) than Si, so the unipolar SiC MOSFET structure is well suited for the implementation of 650, 900 and 1200 V power transistors due to the following main features:

SiC MOSFETs do not have a saturation voltage, unlike Si IGBTs; when paralleling SiC MOSFET chips, the on-resistance can be reduced to $\leq 1\text{--}2$ mOhm;

SiC MOSFETs can conduct in the third quadrant (unlike Si IGBTs) by using a body diode during the dead time (T_{dt} is very short for SiC structures) and then turning on the SiC MOSFET channel in the third quadrant, which gives the same low losses in the reverse conduction state as in the forward conduction state. The combination of using a body diode during the dead time and synchronous rectification eliminates the need for an external antiparallel diode, which reduces the size and cost with minimal impact on efficiency at frequencies up to 50 kHz; using SiC MOSFETs can reduce inverter losses in a typical BEV EPA drive cycle by up to ~78%. The basic technology for developing

low-resistance SiC MOSFET power modules can be scaled from 650-900 V to 1200 V by simply modifying the epitaxial drift zone (blocking layer) and edge regions.

The topology remains the same for all devices in the specified voltage range, ensuring easy integration into power modules (GaN Systems listed on 2021 Deloitte Technology Fast 500). Figure 3 illustrates the traditional method of connecting conductors using ultrasonic welding to the top contact surface using the third generation of SiC MOSFET crystals as an example. This technology can be used in 650, 900 or 1200 V modules with a slight change in the chip topology. 900 V crystals with low channel resistance (10 mOhm for the CPM3-0900-0010A) are already available. They were used in the development of a version of the 900 V modules, the static and dynamic losses of which have already been tested.

Conclusion

The rapid growth of the global automobile market with the gradual predominance of the production of hybrid and electric vehicles guarantees accelerated development and further reduction in the cost of semiconductor devices based on SiC and GaN. Assessing the good market prospects of SiC and GaN products, large foreign companies continue to increase investments in their production, as well as in the acquisition of companies developing these products and materials, including for the start of production of new areas of electronic components that were previously absent in these companies (GaN Systems' power transistor prices drop below \$1, 2021).




Figure 3. Chips Wolfspeed SiC MOSFET

The development of electronic components based on GaN is evolving from the creation of simpler power discrete components (transistors, diodes, etc.) and control microcircuits (drivers, controllers) to more complex integrated solutions with high energy efficiency, including for use in harsh space conditions. Increasing the diameter of SiC wafers to 200 mm, which will happen in 2022 despite technical difficulties, is the main means that will reduce the cost and price of production of silicon carbide ECs.

The first epitaxial GaN wafers on a silicon substrate with a diameter of 300 mm with high homogeneity and low defects, presented on the market in 2021, will further reduce the cost of production and prices of electronic GaN components. Evolution and progress in the industrial development of the technology of vertical GaN-on-GaN and GaN-on-Si structures for active components with low cost in the future will create competition not only for silicon IGBTs, but also for SiC transistors, diodes in the high-voltage range up to 10 kV.

Recently, much attention has been paid to the sintering technology of SiC chips, which allows eliminating the use of welded conductors during assembly. One of the main advantages is an increase in the so-called intermittent service life (IOL), since fatigue processes in welded conductor joints or crystal connections often cause failures. Other potential advantages include better (two-way) cooling, better heat distribution and higher short-circuit resistance (Quantum physics; Semiconductor; HyperPhysics; YOLE Intelligence).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. All About Circuits. (n.d.). *Semiconductor physics introduction*. <https://www.allaboutcircuits.com/video-lectures/semiconductor-physics-introduction/>
2. Bosch. (2021, October 19). *SiC and GaN: A tale of two semiconductors*. <https://www.bosch-presse.de>
3. Encyclopaedia Britannica. (n.d.). *Semiconductor*. <https://www.britannica.com/science/semiconductor>
4. EETimes. (2021, October 19). *SiC and GaN: A tale of two semiconductors*. <https://www.eetimes.com>
5. GaN Systems. (n.d.). *GaN Systems listed on 2021 Deloitte Technology Fast 500*. <https://gansystems.com>
6. HyperPhysics. (n.d.). *Semiconductors*. Georgia State University. <http://hyperphysics.phy-astr.gsu.edu/hbase/Solids/semic.html>
7. Khan Academy. (n.d.). *Quantum physics*. <https://www.khanacademy.org/science/physics/quantum-physics>
8. Power Electronics News. (2021, October 12). *Challenges and future perspectives*. <https://www.powerelectronicsnews.com>
9. PV Education. (n.d.). *Semiconductors*. <https://www.pveducation.org/pvc/drom/pn-junction/semiconductors>
10. Semiconductor Today. (2021, March 10). *GaN Systems' power transistor prices drop below \$1*. <https://www.semiconductor-today.com>
11. Semiconductor Today. (2021, November 18). *GaN Systems raises \$150m in growth capital funding*. <https://www.semiconductor-today.com>
12. YOLE Intelligence. (2021). *Compound semiconductor quarterly market monitor (Q3 2021)*. <https://www.yole.fr>

Tourism Development in Lankaran District and Changes Occurring in Forest Ecosystems

Aynur Heydarova¹ , Alibaghish Malikov² , Tural Huseynov^{3*} 

Abstract. *In the modern era, the tourism sector plays a special role in the global economy while also being important for the development of socio-cultural relations. In recent years, the successful reforms and infrastructure projects implemented in the Republic of Azerbaijan have created a favorable foundation for the sustainable development of tourism. Our country's natural and geographical location, rich cultural and historical heritage, multicultural values, and hospitable people provide broad opportunities for the rapid development of this sector.*

In our republic, the country's future development is determined by the availability of potential economic opportunities and the expansion of international tourism. The activity of numerous foreign companies reflects the integration of entrepreneurs and businesspeople into foreign economic relations and global tourism. Tourism is one of the main directions of development of Azerbaijan's non-oil sector and makes a significant contribution to the process of economic diversification. The social impact of tourism is also evident in increased employment. Thus, new jobs are created in hotel services, public catering, transportation, and other service sectors.

Keywords: *forest, tourism, sector, recreation, dynamics, satellite, infrastructure*

Introduction

Azerbaijan's extensive tourism potential is regarded as a factor that stimulates the development of the tourism industry in its natural and economic regions. However, these opportunities are not being fully and effectively utilized. Therefore, studying the domestic tourism potential of individual regions of the country, including the Lankaran–Astara economic-geographical region examined in this research, identifying ways to use this potential efficiently, forming new tourism services, and applying a wider range of tourism types are highly relevant issues. In Azerbaijan, the number of tourism enterprises has increased by an average of 4.5% annually, employment in the tourism sector by 6%, and the number of foreign citizens visiting the country for tourism purposes by 8.5%. An analysis of the structure of foreign investments directed to the national economy in 2015 shows that 11.3% of total investments were allocated to the non-oil sector, one of the leading branches of which is the tourism sector (Aliyev, 2021).

As a result of the purposeful policy pursued by the President of the Republic of Azerbaijan, Ilham Aliyev, a favorable business environment for the development of tourism has been created in the country, and the share of investments allocated to this sector has increased.

¹ Lankaran State University, PhD in Geography, Lankaran, Azerbaijan

² Lankaran State University, Lankaran, Azerbaijan

³ Lankaran State University, Master's Student, Lankaran, Azerbaijan

*Corresponding author. E-mail: huseynovtural2003@gmail.com

Received: 24 August 2025; Accepted: 14 November 2025; Published online: 25 December 2025

© The Author(s) 2025. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

As noted by the President, tourism has been identified as a priority direction in the development of the non-oil sector, which is reflected in two state programs and the Strategic Roadmap adopted in this field. The main objective of the research is to analyze the current state of tourism in the Lankaran region, identify existing problems and opportunities, and determine future development prospects (Ministry of Ecology and Natural Resources, 2020–2023).

Methods

Lankaran district is a unique region located in the southern part of Azerbaijan, rich in relict and endemic vegetation. Hyrcanian-type forests, which survived the glaciation of the Tertiary period, constitute the main natural wealth of this region. However, over the past decade, the rapid development of tourism, the establishment of new recreation centers, and the increase in anthropogenic pressure have caused significant quantitative and qualitative changes in this fragile ecosystem. The development of tourism in Lankaran has both positive and negative impacts on nature; the positive aspects are aimed at the protection, conservation and promotion of nature (e.g. festivals) thanks to agrotourism (tea, tangerine plantations) and health tourism (thermal waters), while the negative aspects include increased construction, infrastructure expansion, increased waste and pressure on natural resources (water, forests), which risks disrupting the balance of reserves and ecosystems (Mammadov & Khalilov, 2005).

Positive impacts:

- Agrotourism: Tours to tea and tangerine plantations aim to protect and promote nature. For example, tours such as "Green Tea".
- Health tourism: Alongside the use of thermal waters, attention to therapeutic natural resources increases, contributing to their protection.
- Festivals and events: Events such as the "Tea, Citrus, and Rice Growing Festival" enhance the attractiveness of nature and promote eco-tourism.
- Environmental protection: Parallel to tourism development, measures are taken to protect nature, particularly along the Lankaran–Lerik–Astara routes.

Negative impacts (Potential risks):

- Construction and infrastructure: The construction of new hotels and recreation complexes may alter natural landscapes and exert pressure on forest areas.
- Waste management: The increase in tourist flow may lead to waste management problems, negatively affecting rivers and forests.
- Pressure on water resources: The growing water demand of recreation complexes and settlements may put pressure on local water sources.
- Ecosystem changes: The development of mountain and extreme tourism (for example, in forest areas) may interfere with wildlife habitats (UNESCO World Heritage Centre, 2023).

The potential opportunities for tourism development in the region include:

- 1) Increasing the number of foreign tourists;
- 2) Development of infrastructure and modernization of public utilities;
- 3) Prioritizing tourism as a key sector in the development of small and medium-sized enterprises;
- 4) Improving the quality of highways and developing local tourism routes.

Threats to the tourism sector in the region include:

- 1) High intensity of landslide processes, as the foothills and mid-mountain zones of the Talysh Mountains are prone to landslides;
- 2) Fluctuations in the Caspian Sea level, which cause greater damage to the infrastructure of Lankaran and Astara districts;
- 3) Low wage levels;
- 4) Environmental problems in tourism and recreation centers;
- 5) Lack of effective coordination among government institutions;
- 6) The risk of forming a negative image in the long term due to fragmented tourism development;

- 7) Persistence of bureaucratic barriers to tourism development, resulting in weak growth of tourism entrepreneurship;
- 8) Failure to ensure sustainable tourism development due to the prioritization of seasonal and short-term tourism.
- 9) To better understand the impact of tourism, it is essential to analyze the forest cover and infrastructure map of Lankaran. As shown on the map, the main tourism facilities are concentrated around the Khanbulan River reservoir, the Haftoni thermal zone, and along the Lankaran–Lerik highway. These areas are where forest cover is most exposed to fragmentation (WWF – World Wide Fund for Nature, 2023).

Results and Discussion

Statistical Indicators and Dynamics:

There is a direct correlation between tourist flow and the degradation of forest areas.


Table 1.
Condition of forest cover around tourism centers

Year	Numbers of tourism	Canopy cover (0-1 scale)	Soil compaction (kg/cm ²)
2015	45,000	0.82	1.4
2020	78,000	0.76	2.8
2024	115,000	0.69	4.1

The data show that soil compaction has increased by approximately three times over the past nine years. This leads to the destruction of understory vegetation and young saplings. The dominant forest species of Lankaran—ironwood (*Parrotia persica*) and chestnut-leaved oak (*Quercus castaneifolia*)—are particularly sensitive to noise and light pollution generated by tourism facilities (Ministry of Ecology and Natural Resources, 2020–2023).

Comparison with International Experience :

Compared to Germany’s Schwarzwald (Black Forest) region, sustainable tourism practices are weakly implemented in Lankaran. In Germany, special suspension bridges and wooden walkways are constructed for tourists to minimize direct contact with the soil. The topographic structure of Lankaran is a key factor determining the degree of tourism penetration into forest areas. The relief map clearly shows that forest massifs extend from the Lankaran lowland (below sea level) up to the ridges of the Talysh Mountains. This is a physical map illustrating relief, river networks, and elevation zones (Map 1).



Map 1. Physical–topographic map of the Lankaran–Astara region (2022–2024)
(Based on satellite imagery)

Hydrographic Network: The valleys of the Lankaran River, Basharu River, and Girdanichay visible on the map are the areas where tourism is developing most actively. The topographic map demonstrates that anthropogenic pressure is concentrated mainly in forested areas along river valley slopes.

Slope Gradient: Tourism roads constructed in areas with slope gradients of 20–30° accelerate erosion processes. The topographic map indicates which forest sections—particularly steep slopes—should be designated as “restricted zones” for tourism.

The physical–topographic map of Lankaran (relief and hydrography) clearly reflects the elevation zones in which forests are located and the relief forms (river valleys, slopes) where tourism is geographically concentrated. Analysis of the topographic map shows that tourism infrastructure is mainly concentrated along river valleys at elevations of 0–400 meters above sea level. These areas also represent the most productive and biologically rich forest zones, encompassing lowland and foothill belts. While the complexity of the relief prevents the large-scale expansion of tourism to higher elevations (above 600 meters), it intensifies ecological pressure within the valleys (UNESCO World Heritage Centre, 2023).

This map illustrates the actual health condition of the forest. Dark green colors indicate the densest and most intact forest areas, while lighter shades represent thinned zones affected by tourism and other anthropogenic factors. Overall, the topographic analysis confirms that tourism infrastructure in Lankaran is primarily concentrated along river valleys (Lankaran River and Basharu River) at elevations of 0–400 meters above sea level. As shown on the map, although mountainous relief naturally restricts mass tourism from expanding into higher zones, it transforms forest belts in river valleys into focal points of anthropogenic pressure.

Modern Forest Cover and Vegetation Density of Lankaran

The modern forest map (based on Sentinel-2 satellite imagery) shows that the forest massifs of the district differ sharply in both species composition and canopy density with increasing elevation above sea level (Map 2).



Map 2. Joint satellite imagery map for 2024 by “Azercosmos” OJSC and the monitoring group of the Ministry of Ecology and Natural Resources


Dark Green Zones: Areas with high humidity and minimal human intervention. These mainly include the core parts of Hirkan National Park (reserve zones) (State Tourism Agency of Azerbaijan, 2023).
Light Green and Yellowish Zones: Areas close to tourism zones where forest canopy density has decreased. These include forest road edges and newly established cottage-type recreation areas.

Zone A (Coastal and Lowland Forests): The area extending from the Caspian Sea coastline to the main highway. Tourism facilities are most densely concentrated here, and forest cover is highly fragmented.

Zone B (Foothill and Lower Mountain Belt – 200–600 m): This zone includes the main entrances to Hirkan National Park, the surroundings of the Khanbulan River reservoir, and the “Istisu” area. It is considered a “red zone” where tourism causes the greatest physical damage to forests.

Zone C (High Mountain and Reserve Zone – above 600 m): A zone with restricted tourist access, where endemic species—such as the healthiest populations of boxwood and ironwood—are protected (Mammadov, 2007).

The analysis of maps of recreation centers and tourism routes in the Lankaran–Astara economic-geographical region, together with NDVI (Normalized Difference Vegetation Index) values obtained from satellite imagery, confirms that forest canopy density within a 500–800 meter radius of main highways and tourism centers has decreased by 12–15% over the past decade. This clearly demonstrates the fragmentation of the internal forest ecosystem. Satellite imagery enables the measurement of the “greenness” level of vegetation, providing an objective assessment of vegetation health and density (Map 3) (WWF – World Wide Fund for Nature, 2023).



Map 3. Map of Recreation Centers and Tourism Routes in the Lankaran–Astara Economic-Geographical Region

Based on the example of Lankaran district, GIS studies conducted over the past 10 years show that within a 500-meter radius of tourism centers, the NDVI value has decreased on average from 0.85 to 0.70. This indicates forest thinning and a decline in photosynthetic capacity

- Khanbulan Reservoir: Classified as an « ecological stress zone » due to high tourist density
- Lankaran–Lerik Road: Areas where roadside cafés cut through forest belts in a corridor-like pattern.
- Haftoni settlement surroundings: A zone where medical tourism (thermal waters) has altered groundwater levels, affecting forest moisture conditions.
- The development of tourism primarily requires the construction of road and hotel infrastructure. This process has led to the following changes in forests:
- Road construction: New roads built to remote mountain villages of Lerik and Astara have caused fragmentation of forest massifs, affecting migration routes of certain animal species.
- Recreation centers: Cottages and restaurants built on forest fund lands have resulted in a certain reduction in tree density. However, in recent years, the application of “green construction” principles have been observed, integrating infrastructure into forest areas without tree cutting.

Tourism development does not result solely in negative impacts; it has also created new opportunities for forest conservation:

- Ecotourism routes: Tourists now visit forests not only for picnics but also for birdwatching and hiking, increasing interest in preserving forests in their natural state
- Role of national parks: Enhancing the tourism potential of Hirkan National Park has strengthened state-level oversight. Funding allocated for forest protection and related services has increased.
- Local community income: As tourism develops, local residents become more interested in protecting forests as a “tourism product” rather than cutting trees for firewood (Yusifov & HajiyeV, 2004).

Conclusion

- Although tourism development in Lankaran is oriented toward nature conservation, without continuous planning, ecological monitoring, and the application of sustainable tourism principles, construction and human flow may negatively impact the natural environment.
- Satellite imagery has provided precise evidence of forest thinning associated with tourism development.
- Tourism facilities should be constructed only along forest edges within designated buffer zones.
- A portion of taxes collected from tourism facilities should be allocated to forest restoration.
- Organizing tourist routes based on the region’s existing tourism potential allows for attracting more visitors. Therefore, offering ecological, national heritage, and educational tourism routes to visitors in the Lankaran–Astara economic-geographical region is essential.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Aliyev, T. (2021). Regional aspects of tourism development in Azerbaijan. *Journal of Economics*.
2. Hall, C. M., & Page, S. (2014). *The Geography of Tourism and Recreation*. Routledge.
3. Ministry of Ecology and Natural Resources. (2020–2023). *Annual reports on the ecological condition of Hirkan National Park*.
4. Mammadov, G. Sh., & Khalilov, M. Y. (2002). *Forests of Azerbaijan*. Elm.
5. Mammadov, G. Sh., & Khalilov, M. Y. (2005). *Ecology and Environmental Protection*. Elm.
6. Mammadov, G. Sh. (2007). *Land Resources of Azerbaijan*.
7. Rzayev, A. (2019). *Regional tourism resources and development directions*. Adiloghlu.
8. State Statistical Committee of the Republic of Azerbaijan. (2024). *Environment of Azerbaijan – 2023: Statistical Yearbook*.
9. State Tourism Agency of Azerbaijan. (2023). *Regional Tourism Development Strategy 2023–2026*.
10. Sharples, R. (2018). *Tourism, Sustainable Development and the Environment*. Routledge.
11. UNESCO World Heritage Centre. (2023). «Hyrkanian Forests (Azerbaijan/Iran)» nomination dossier.
12. UNWTO (World Tourism Organization) – *Official reports and analytical materials*.
13. WWF (World Wide Fund for Nature). (2023). “Caucasus Ecoregion Strategy 2020–2025” report.
14. Yusifov, E. F., & HajiyeV, V. C. (2004). *Hirkan Biosphere Reserve*. El-Alliance.

CONTENTS

Nurgun Mehbaliyeva, Narmina Abdullayeva, Nigar Huseynova Metabolism of the Bilirubin and its Biochemistry Role	4
Ayten Agayeva, Rahila Farmanli Study of Diagnostic Methods and Control Measures For Listerosis Disease Among Big-Horned Animals in the Ganja-Gazakh Zone.....	10
Gunay Dadashzade The Possibilities of Using the Nature of Semiconductors in Practical Life	14
Aynur Heydarova, Alibaghish Malikov, Tural Huseynov Tourism Development in Lankaran District and Changes Occurring in Forest Ecosystems	20

Editorial address

AZ1073, Baku,
Matbuat Avenue, 529,
“Azerbaijan” Publishing House, 6th floor
Phone: +994 99 805 67 68
+994 99 808 67 68
e-mail: nature.science2000@aem.az

Signed: 19.12.2025
Online publication: 25.12.2025
Paper printing: 10.01.2026
Format: 60/84, 1/8
Stock issuance: 2,75 p.s..
Order: 124

It has been published in the printing house
“ZANGAZURDA”
Address: Baku city, Matbuat Avenue, 529,
“Azerbaijan” Publishing House, 6th floor
Phone: +994 12 510 63 99
e-mail: zengezurda1868@mail.ru

